

Chapter 7. Conclusions

In the first part of this Ph.D. Thesis we have studied the optical correlator. We have performed the optimization of the modulation response for both the scene SLM and the filter SLM. We have also proposed an alignment procedure for the correlator. In the second part of the Thesis we have proposed and studied the description of color images by means of three dimensional functions, and we have defined the Fourier transform and correlation for this kind of signals. Finally, in the third part of the Thesis we have implemented the three dimensional correlation of color images in the optical processor. The obtained results lead to the following conclusions:

- ✧ A formalism based on the description of Jones vectors by means of Dirac brackets has been introduced. The proposed formalism permits to separate the wave information from the polarization state information carried out by a Jones vector. The operators corresponding to different polarizing elements have been given for the proposed formalism.

- ✧ The introduced formalism has been used to prove that the amplitude and phase modulation of spatial light modulators depend only on the polarization state generated at the input side of the panel and the polarization state detected at the output of the panel. The expression of the modulation as a function of the four free parameters that determine these polarization states has been given.
- ✧ The representation of polarization states in Poincaré's sphere has been used to establish the relation between the polarization states and the polarization state generator and detector used to generate or detect these states. The retardance requirements of the retarder plates used in the optical systems have been determined.
- ✧ The developed formalism has been applied to optimize the modulation of the liquid crystal spatial light modulators used in the optical correlator. We have defined a merit function to evaluate the quality of the modulation response of the spatial light modulator as amplitude only operation curve. We have also defined a merit function for the phase only modulation regime.
- ✧ An iterative algorithm for the optimization of the merit functions has been proposed. This algorithm is capable of obtaining the absolute maximum for the merit function, and not only the local maxima. We have used the proposed algorithm to obtain the polarization states at the input and the output of the

liquid crystal panel that optimize the modulation of the spatial light modulators of the correlator. We have obtained the solutions for amplitude only modulation for the scene SLM, and the solutions of phase only modulation for both the scene and the filter SLMs.

- ✧ The configuration of the polarizing elements that generate and detect the optimal states have been obtained. These configurations have been implemented in the optical correlator and the corresponding operation curves have been calibrated.
- ✧ A procedure based on simple scene and filter tests have been proposed to perform the alignment of the optical convergent correlator for either amplitude or phase encoded scene. All the alignment sequence is based on the observation of the reconstructed image of the scene in the correlation plane and the interpretation of the effects of the filtering.
- ✧ The tests we have proposed permit the focusing of the diffraction pattern of the scene on the filter plane, as well as the centering of the filter, the alignment of the azimuth between the scene and the filter SLMs, and the scaling of the Fourier spectrum of the scene on the filter plane.
- ✧ The proposed tests constitute a series of objective criteria for the alignment of the correlator, in the sense that experimental observations are interpreted in accordance to a simple frequency filtering theory. For all the tests we have presented, experimental results that demonstrate the correct alignment of

the optical correlator have been given. We have shown that the proposed tests constitute a good method to solve the main drawback (the alignment requirements) of the Vander Lugt type correlators.

In the second part of this Ph.D. Thesis we have introduced the description of color images, or hyper-spectral images, by means of three dimensional functions in which the third coordinate contains the color information. We have analyzed the color coordinate as a sample of the spectral distribution in each one of the points of the color image.

- ✧ The color Fourier transform, that is, the Fourier transform along the color axis has been defined. We have also defined the three dimensional Fourier transform for three dimensional functions that describe color images.
- ✧ We have shown that each one of the channels of the color Fourier transform represents an orthogonal projection onto one subspace of the corresponding color space. The subspaces corresponding to the different channels of the color Fourier spectrum are orthogonal each other.
- ✧ In particular, the projections corresponding to the channels of the color Fourier spectrum for the case of RGB color images have been determined. We have determined that the $m=0$ channel of the color Fourier spectrum contains the information of the intensity of the color, and that the chromaticity information is contained in the $m=1$ and $m=-1$ channels of the

color Fourier spectrum. The saturation is the ratio of the magnitude of the $m=1$ channel to the $m=0$ channel, and the hue is the argument of the $m=1$ channel.

- ✧ The main properties of the Fourier transform for the case of three dimensional functions that describe color images have been given. We have shown that the three dimensional spectrum of color images contains the interference between the objects in different channels of the image.
- ✧ The three dimensional correlation of three dimensional functions has been proposed as a valid technique for color pattern recognition. We have established the conditions for the three dimensional correlation to be useful for color pattern recognition. In particular we have demonstrated that, for recognition purposes, the three dimensional correlation function must be evaluated in the origin of the color axis because the color of the objects to recognize is not invariant for shifts along the color axis of the three dimensional functions that describe them.
- ✧ An interpretation of the filtering operations of the color Fourier spectrum as linear transformations of the color space have been given. We have provided a classification of the linear filtering operations. The high-pass and low-pass filters of the color Fourier spectrum have been interpreted, respectively, as dilatations and contractions of the color space along the axis normal to the $1/3$ -intensity plane. The effect of the high pass

filter to the color Fourier spectrum involves an enhancement of the saturation of the colors in the image described by the filtered image, keeping unaltered the hue of the colors. The effect of the low-pass filter applied to the color Fourier spectrum of an image involves a reduction of the saturation of the colors in the reconstructed color image, keeping unaltered the hue of the colors. We have demonstrated that the only phase transformation of the color Fourier spectrum of an image is the application of a linear phase. The linear phase filtering involves a rotation of the color space about the normal to the 1/3-intensity plane. This rotation involves a modification of the hue of the colors of the filtered image, but the intensity and saturation are not altered.

- ✧ The interpretation for some non-linear operations such as the whitening of the Fourier spectrum has been given. We have demonstrated that the whitening of the color Fourier spectrum of a three dimensional function that describes a color image involves a normalization of the intensity of the colors and a maximization of the saturation of the colors of the image. We show that most of the colors resulting from the whitening of the color Fourier spectrum have negative components. This is interpreted as the amount of the corresponding primary color that must be added to the saturated color to obtain a non-saturated color that can be composed by addition of the remaining primary colors.

- ✧ The color-wise correlation of an image has been defined as the correlation of the three dimensional function that describes the image along the color axis, considering the addition of all the possible cyclic translations of the image along the color axis.
- ✧ The interpretation of the classic matched filter for color-wise correlation is given. We show that the color-wise correlation, evaluated at the origin of the color axis gives the scalar product between the vectors that point to the colors in the color space. Therefore the color-wise autocorrelation is maximum for the target color in comparison to the cross-correlations of the target color with any other color with equal or lower color energy.
- ✧ The phase only filter has been interpreted. We have shown that the color wise correlation using phase only filter represents a change between different orthogonal bases of the color space. In addition, because the whitening of the color Fourier spectrum is equivalent to a normalization of the intensity and the saturation of the colors of the image, we have provided an alternative path to perform the color wise correlation using this filter.
- ✧ The interpretation for the pure phase color-wise correlation is given. We have shown that in this case only the hue of the colors is taken into account for the recognition, and therefore the colors are classified by their hue.
- ✧ We have also considered whitening operations on the three dimensional Fourier transform of color images, and its implications on color pattern recognition. When the three

dimensional Fourier spectrum of an image is whitened, the filtered image presents the typical edge enhancement of the colors, and the colors are also normalized (but the saturation is not exactly maximized). In this case the three dimensional autocorrelation is a sharp peak in the origin, and null elsewhere. This indicates both that the channels are decorrelated each other, and that the channels are also spatially decorrelated.

- ✧ The discrimination capability for three dimensional correlation involving different whitening operations for objects with similar shape and different color distribution has been compared. The whitening of the scene color Fourier spectrum is also proposed because it can be performed in the direct domain by means of normalizing the intensity and the saturation of the colors of the images.

In the third part of the Thesis, we have proposed a technique to encode the three dimensional functions that describe color images in two dimensional functions. The technique allows us to obtain the three dimensional Fourier transform and the three dimensional correlation.

- ✧ The proof that the proposed encoding can be used to obtain the three dimensional Fourier spectrum of the encoded image has been given. We have shown that the obtained Fourier transform corresponds to a non-orthogonal sample of the three dimensional Fourier spectrum of the three dimensional image.

- ✧ An interpretation of the encoding in terms of shifted carriers has been provided. This allows us to determine the sampling requirements for the encoded function, that leads to the orthogonal sampling of the three dimensional Fourier spectrum.
- ✧ We have also proposed a variant of the encoding technique in which one of the spatial coordinates is encoded on the color coordinate. We have also demonstrated that in this case the channels of the Fourier spectrum of the three dimensional image are interlaced.
- ✧ The proof that also the three dimensional correlation can be obtained by means of two dimensional operations by the proposed encoding technique has been given. We demonstrate that the $n=0$ channel is encoded exactly. In addition, for the other channels the error is very small if the values of two consecutive pixels are not very different.
- ✧ The implementation in the optical correlator of the proposed encoding technique has been performed. The bandwidth limitations of the optical system and the alignment requirements have been taken into account. Aside from the alignment procedure for the optical correlator, we have developed a Moiré technique to control the scale of the correlation plane with the camera CCD array, which is necessary for the proposed encoding.

- ✧ We have implemented three variants of the encoding technique: considering full bandwidth color encoding, half bandwidth color encoding and spatial encoding. The three dimensional correlation for color pattern recognition has been performed in the optical correlator using the three variants. We have obtained excellent results for both the three dimensional phase only filter and the color spectrum whitened scene, using a three dimensional phase only filter.

✧

References

- [Abookasis2001]** D. Abookasis, O. Arazi, J. Rosen, B. Javidi, "Secure optical systems based on a joint transform correlator with significant output images", *Opt. Eng.* **40**, pg.1584-1589 (2001).
- [Ahouzi94]** E. Ahouzi, J. Campos, M.J. Yzuel, "Phase-only filter with improved discrimination ", *Opt. Lett.* **19**, pg.1340-1342 (1994).
- [Ahouzi98]** E. Ahouzi, J. Campos, M.J. Yzuel, "Binary amplitude phase-only filter with high multiobject discrimination capability", *Opt. Eng.* **37**, pg.2351-2358 (1998).
- [Alam2001]** M.S. Alam, C.N. Wai, "Color pattern recognition using fringe adjusted joint transform correlator", *Opt. Eng.* **40**, pg.2407-2413 (2001).

- [Attaleb99]** A. Attaleb, E. Ahouzi, J. Campos, M.J. Yzuel, "Optical color pattern recognition with high discrimination capability using binary amplitude phase-only filter", *Opt. Rev.* **6**, pg.42-48 (1999).
- [Badiqué88A]** E. Badiqué, N. Ohyama, T. Honda, J. Tsujiuchi, "Color image correlation for spatial/spectral recognition and increased selectivity", *Opt. Commun.* **68**, pg.91-96 (1988).
- [Badiqué88B]** E. Badiqué, Y. Komiya, N. Ohyama, T. Honda, J. Tsujiuchi, "Use of color image correlation in the retrieval of gastric surface topography by endoscopic stereopair matching.", *Appl. Opt.* **27**, pg.941-948 (1988).
- [Barbé99]** J. Barbé, J. Campos, "Image segmentation with a white light optical correlator", *Proc SPIE Vol. 3749*, , pg.775-776 (1999).
- [Bhattacharya2002]** N. Bhattacharya, H.B. van Linden van den Heuvell, R.J.C. Spreeuw, "Implementation of quantum search algorithm using classical Fourier optics", *Phys Rev. Lett.* **88**, pg.137901- (2002).
- [Burckhardt70]** C.B. Burckhardt, "A simplification of Lee's method of generating holograms by computer", *Appl. Opt.* **9**, pg.1949- (1970).
- [Burr99]** G.W. Burr, S. Kobras, H. Hanssen, H. Coufal, "Content-addressable data storage by use of volume holograms", *Appl. Opt.* **38**, pg.6779-6784 (1999).

- [Busarow2000]** P.L. Busarow, A. Mahalanobis, "Utilization of an optical correlator for automatic target recognition against ballistic missile defense targets", *Proc. SPIE Vol. 4043, D.P. Casasent, T.H. Chao, Eds.*, pg.140-148 (2000).
- [Cai96]** L. Cai, Y. Jin, S. Zhou, P. Yeh, N. Marzwell, H. Liu, "Translational sensitivity adjustable compact optical correlator and its application for fingerprint recognition", *Opt. Eng.* **35**, pg.415-422 (1996).
- [Campos91]** J. Campos, M.S. Millán, M.J. Yzuel, C. Ferreira, "Colour invariant character recognition and character-background colour identification by multichannel matched filter", *Proc SPIE, Vol. 1564* , pg.189-198 (1991).
- [Case79]** S.K. Case, "Pattern recognition with wavelength-multiplexed filters", *Appl. Opt.* **18**, pg.1890-1894 (1979).
- [Caulfield97]** H.J. Caulfield, I. Moreno, J. Campos, M.J. Yzuel, "Coherent recognition of colored patterns", *Opt. Commun.* **133**, pg.77-81 (1997).
- [Chang97]** S. Chang, S.A. Boothroyd, J. Chrostowski, "Partial rotation-invariant pattern matching and face recognition with a joint transform correlator", *Appl. Opt.* **36**, pg.2380-2388 (1997).

- [Chang96]** S. Chang, S.A. Boothroyd, P. Palacharla, S. Pachanathan, "Rotation invariant pattern recognition using a joint transform correlator.", *Opt. Commun.* **127**, pg.107-116 (1996).
- [Chang2002]** H.T. Chang, W.C. Lu, C.J. Kuo, "Multiple-phase retrieval for optical security systems by use of random phase encoding", *Appl. Opt.* **41**, pg.4825-4834 (2002).
- [Chao2002]** T.-S. Chao, H. Zhou, G.F. Reyes, , "Spacecraft navigation using a grayscale optical correlator", *Proc. SPIE Vol. 4734*, D.P. Casasent, T.-H. Chao, Eds., pg.108-113 (2002).
- [Cohn94]** R.W. Cohn, M. Liang, "Approximating fully complex spatial modulation with pseudorandom phase-only modulation ", *Appl. Opt.* **33**, pg.4406-4415 (1994).
- [Cohn98]** R.W. Cohn, "Pseudorandom encoding of complex-valued functions onto amplitude-coupled phase modulators", *J. Opt. Soc. Am. A* **15**, pg.868-883 (1998).
- [Cohn99]** R.W. Cohn, M. Duelli, "Ternary pseudorandom encoding of Fourier transform holograms", *J. Opt. Soc. Am. A* **16**, pg.71-84 (1999).
- [Cook98]** N.J. Cook, A. Carnicer, S. Vallmitjana, I. Juvells, C.M. Cartwright, W.A. Gillespie, "implementation of a photorefractive binary joint transform correlator", *J. Opt. Soc. Am. B* **15**, pg.1977-1984 (1998).

- [Corbalán2002]** M. Corbalán, M.S. Millán, M.J. Yzuel , "Color pattern recognition with CIELAB coordinates", *Opt. Eng.* **41**, pg.130-138 (2002).
- [Coy96]** J.A. Coy, M. Zaldarriaga, D.F. Grosz, O.E. Martínez, "Characterization of a liquid crystal television as a programmable spatial light modulator", *Opt. Eng.* **35**, pg.15-19 (1996).
- [Davis89]** J. A. Davis, M.A. Waring, G.W. Bach, R.A. Lilly, D.M. Cottrell, "Compact optical correlator design", *Appl. Opt.* **28**, pg.10-11 (1989).
- [Davis98]** J.A. Davis, I. Moreno, P. Tsai, "Polarization eigenstates for twisted-nematic liquid-crystal displays", *Appl. Opt.* **37**, pg.937-945 (1998).
- [Davis99A]** J.A. Davis, D.M. Cottrell, J. Campos, M.J. Yzuel, I. Moreno, "Encoding amplitude information onto phase-only filters", *Appl. Opt.* **38**, pg.5004-5013 (1999).
- [Davis99B]** J.A. Davis, P. Tsai, K.G. D'Nelly, I. Moreno, "Simple technique for determining the extraordinary axis direction for twisted nematic liquid crystal spatial light modulators", *Opt. Eng.* **38**, pg.929-932 (1999).
- [Davis99C]** J.A. Davis, J.C. Escalera, J. Campos, A. Márquez, M.J. Yzuel, C. Iemmi, "Programmable axial apodizing and hyperresolving amplitude filters with a liquid-crystal spatial light modulator", *Opt. Lett.* **24**, pg.628-639 (1999).

[Davis99D] J.A. Davis, D.B. Allison, K.G. D'Nelly, M.L. Wilson, I. Moreno, "Ambiguities in measuring the physical parameters for twisted-nematic liquid crystal spatial light modulators", *Opt. Eng.* **38**, pg.705-709 (1999).

[Davis2002] J.A. Davis, M.D. Mowak, "Selective edge enhancement of images with an acousto-optic light modulator", *Appl. Opt.* **41**, pg.4835-4839 (2002).

[Duelli97] M. Duelli, A.R. Pourzand, N. Collings, R. Dndliker, "Pure phase correlator with photorefractive filter memory", *Opt. Lett.* **22**, pg.87-89 (1997).

[Duelli99] M. Duelli, R.W. Cohn, "Pseudorandom encoding for real-valued ternary spatial light modulators", *Appl. Opt.* **38**, pg.3804-3809 (1999).

[Efron95] U. Efron, *Spatial Light Modulator technology: materials, devices and applications* , Marcel Dekker, New York (1995).

[Esteve-Taboada2000] J.J.Esteve-Taboada, J. García, C. Ferreira, "Extended scale-invariant pattern recognition with white-light illumination ", *Appl. Opt.* **39**, pg.1268-1271 (2000).

- [Fernández-Pousa2000]** C.R. Fernández-Pousa, I. Moreno, N. Bennis, C. Gómez-Reino, "Generalized formulation and symmetry properties of reciprocal nonabsorbing polarization devices: application to liquid-crystal displays", *J. Opt. Soc. Am. A* **17**, pg.2074-2080 (2000).
- [Fernández-Pousa2001]** C.R. Fernández-Pousa, I. Moreno, J.A. Davis, J. Adachi, "Polarizing diffraction-grating triplicators", *Opt. Lett.* **26**, pg.1651-1653 (2001).
- [Ferreira92]** C. Ferreira, M.S. Millán, M.J. Yzuel, J. Campos, "Experimental results in color pattern recognition by multichannel matched filtering", *Opt. Eng.* **31**, pg.2231-2238 (1992).
- [Florence89]** J.M. Florence, "Joint-transform correlator systems using deformable-mirror spatial light modulators", *Opt. Lett.* **14**, pg.341-343 (1989).
- [Gabor48]** D. Gabor, "A new microscopic principle", *Nature* **161**, pg.777-778 (1948).
- [Gabor49]** D. Gabor, "Microscopy by reconstructed wave-fronts ", *Proc. Royal Soc. Biol.* **A197**, pg.454-487 (1949).
- [González92]** RC. González, R.E. Woods, *Digital Image Processing* , Addison-Wesley (1992).

- [Griffin94]** R.D. Griffin, J.N. Lee, "Acousto-optical wideband correlator system: design, implementation and evaluation", *Appl. Opt.* **33**, pg.6774- (1994).
- [Gualdrón93]** O. Gualdrón, H.H. Arsenault, "Phase-derived circular harmonic filter", *Opt. Commun.* **104**, pg.32-34 (1993).
- [Gualdrón96]** O. Gualdrón, H.H. Arsenault, "Optimum rotation-invariant filter for disjoint-noise scenes", *Appl. Opt.* **35**, pg.2507-2513 (1996).
- [Guillaume97]** M. Guillaume, P. Réfrégier, J. Campos, V. Lashin, "Detection theory approach to multichannel pattern location", *Opt. Lett.* **22**, pg.1887-1889 (1997).
- [Guillaume98]** M. Guillaume, P. Melon, P. Réfrégier, A. Llebaria, "Maximum-likelihood estimation of an astronomical image from a sequence at low photon levels", *J. Opt. Soc. Am. A* **15**, pg.2841-2848 (1998).
- [Hao2002]** Y.Hao, W. Liu, S. Lu, Y. Zhang,, "Applications of small optical correlator in traffic", *Proc. SPIE, Vol. 4921, Y. Zhang, W. Liu, H.M. Pollicove, Eds. ,* pg.71-77 (2002).
- [Hassebrook96]** L.G. Hassebrook, M.E. Lhamon, R. Daley, R.W. Cohn, M. Liang, "Random phase encoding of composite fully complex filters", *Opt. Lett.* **21**, pg.272-274 (1996).

- [Hester80]** C.F. Hester, D. Casasent, "Multivariant technique for multiclass pattern recognition", *Appl. Opt.* **19**, pg.1758-1761 (1980).
- [Horner84]** J.L. Horner, P.D. Gianino, "Phase-only matched filtering", *Appl. Opt.* **23**, pg.812-816 (1984).
- [Hsieh2002]** M.-L. Hsieh, K.Y. Hsu, H. Zhai, "Color image recognition by use of a joint transform correlator of three liquid crystal televisions", *Appl. Opt.* **41**, pg.1500-1504 (2002).
- [Hsu82]** Y.-N. Hsu, H.H. Arsenault, "Optical pattern recognition using circular harmonic expansion", *Appl. Opt.* **21**, pg.4016-4019 (1982).
- [Huard94]** S. Huard, *Polarisation de la lumière*, Masson, Paris (1994).
- [Iwaki90]** T. Iwaki, Y. Mistuoka, "Optical pattern recognition of letters by a joint-transform correlator using a ferroelectric liquid crystal spatial light modulator", *Opt. Lett.* **15**, pg.1218-1220 (1990).
- [Javidi89A]** B. Javidi, "Nonlinear joint power spectrum based optical correlation", *Appl. Opt.* **28**, pg.2358-2367 (1989).
- [Javidi89B]** B. Javidi, "Synthetic discriminant function-based binary nonlinear optical correlator", *Appl. Opt.* **28**, pg.2490-2493 (1989).

- [Javidi94]** B. Javidi, J.L. Horner, "Optical pattern recognition for validation and security verification ", *Opt. Eng.* **33**, pg.1752-1756 (1994).
- [Javidi96]** B. Javidi, G. Zhang, J. Li, "Experimental demonstration of the random phase encoding for image encryption and security verification ", *Opt. Eng.* **35**, pg.2506-2512 (1996).
- [Javidi98]** B. Javidi, E. Ahouzi, "Optical security system with Fourier plane encoding", *Appl. Opt.* **37**, pg.6247-6255 (1998).
- [Javidi2000]** B. Javidi, T. Nomura, "Securing information by use of digital holography", *Opt.Lett* **25**, pg.28-30 (2000).
- [Jerrard54]** H.G. Jerrard, "Transmission of light through birefringent and optically active media: the Poincaré sphere", *J. Opt. Soc. Am.* **44**, pg.634-640 (1954).
- [Jones41]** R.C. Jones, "A new calculus for the treatment of optical systems. Description and discussion of the calculus", *J. Opt. Soc. Am.* **31**, pg.488-493 (1941).
- [Juday93]** R.D. Juday, "Optimal realizable filters and the minimum Euclidean distance principle", *Appl. Opt.* **32**, pg.5100-5111 (1993).
- [Juvells92]** I. Juvells, S. Vallmitjana, S. Bosch, "Analysis of a scale tunable telephoto lens and its use in optical correlation", *J. Modern Optics* **39**, pg.1107-1115 (1992).

- [Keyes2001]** R.W. Keyes, "Fundamental Limits of Silicon Technology", *Proc. IEEE* **89**, pg.227-239 (2001).
- [Kim2002]** J.-J. Kim, J.-H. Choi, E.-S. Kim, "Robust information hiding and extraction system based on complex phase code and optical correlator", *Proc. SPIE Vol. 4677, R.L van Renesse, Ed.*, pg.357-365 (2002).
- [Kirsch90]** J.C. Kirsch, D.A. Gregory, "Video rate optical correlation using a magneto-optic spatial light modulator", *Opt. Eng.* **29**, pg.1122-1128 (1990).
- [Kober97]** V. Kober, V. Lashin, I. Moreno, J. Campos, L.P. Yaroslavsky, M.J. Yzuel, "Color component transformation for optical pattern recognition", *J. Opt. Soc. Am. A* **14**, pg.2656-2669 (1997).
- [Leclerc82]** L. Leclerc, Y. Sheng, H.H. Arsenault, "Rotation invariant phase-only and binary phase-only correlation", *Appl. Opt.* **28**, pg.1251-1256 (1982).
- [Ledesma98]** S. Ledesma, C. Iemmi, J. Campos, M.J. Yzuel, "Joint transform architecture with a single LCTV operatin in phase-mostly mode", *Opt. Commun.* **151**, pg.101-109 (1998).
- [Lee70]** W.H. Lee, "Sampled Fourier transform hoogram generated by computer", *Appl. Opt.* **9**, pg.639-643 (1970).

- [Leith2000]** E.N. Leith, "The evolution of information optics ", *IEEE Journal of selected topics in quantum electronics* **6**, pg.1297-1304 (2000).
- [Liu85]** H.K. Liu, J.A. Davis, R.A. Lilly, "Optical data processing properties of a liquid-crystal television spatial light modulator", *Opt. Lett.* **10**, pg.635-637 (1985).
- [Lohman67]** A. W. Lohman, D.P. Paris, "Binary Fraunhofer holograms, generated by computer", *Appl. Opt.* **6**, pg.1739- (1967).
- [Lu90]** K. Lu, B.E.A. Saleh, "Theory and design of the liquid crystal TV as an optical spatial phase modulation", *Opt. Eng.* **29**, pg.240-246 (1990).
- [Mahalanobis87]** A. Mahalanobis, B.V.K Vijaya Kumar, D. Casasent, "Minimum average correlation energy filters", *Appl. Opt.* **31**, pg.1823-1833 (1987).
- [Márquez2000]** A. Márquez, J. Campos, M.J. Yzuel, I. Moreno, J. Davis, C. Iemmi, A. Moreno, A. Robert, "Characterization of edge effects in twisted nematic liquid crystal displays", *Opt. Eng.* **39**, pg.3301-3307 (2000).
- [Márquez2001A]** A. Márquez, C. Iemmi, I. Moreno, J.A. Davis, J. Campos, M.J. Yzuel, "Quantitative predictions of the modulation behavior of twisted nematic liquid crystal displays based on a simple physical model", *Opt. Eng.* **40**, pg.2558-2564 (2001).

- [Márquez2001B]** A. Márquez, C. Iemmi, J.C. Escalera, J. Campos, S.Ledesma, J.A. Davis, M.J. Yzuel, "Amplitude apodizers encoded onto Fresnel lenses implemented on a phase-only spatial light modulator", *Appl. Opt.* **40**, pg.2316-2322 (2001).
- [McAulay2000]** A.D. McAulay, "Optical correlator for improving images distorted by atmospheric turbulence", *Proc. SPIE Vol. 4046, D.R. Pape, Ed.*, pg.41-46 (2000).
- [McCabe2000]** A. McCabe, T. Caelli, G. West, A. Reeves, "Theory of spatiochromatic image encoding and feature extraction", *J. Opt. Soc. Am. A* **17**, pg.1744-1754 (2000).
- [Millán89]** M.S. Millán, J. Campos, C. Ferreira, M.J. Yzuel, "Matched filter and phase-only filter performance in color image recognition", *Opt. Commun.* **73**, pg.277-284 (1989).
- [Millán92]** M.S. Millán, M.J. Yzuel, J. Campos, C. Ferreira, "Different strategies in optical recognition of polychromatic images", *Appl. Opt.* **31**, pg.2560-2567 (1992).
- [Millán93]** M.S. Millán, M.J. Yzuel, "High pass matched filtering for optical character recognition with character-background contrast invariance", *Proc SPIE Vol. 1983* , pg.421-422 (1993).
- [Millán95]** M.S. Millán, M. Corbalán, J. Romero, M.J. Yzuel, "Optical pattern recognition based on color vision models", *Opt. Lett.* **20**, pg.1722-1724 (1995).

- [Mogensen2000]** P.C. Mogensen, J. Gückstad, "Phase-only optical encryption", *Opt. Lett.* **25**, pg.566-568 (2000).
- [Montes-Usátegui97]** M. Montes-Usategui, S.E. Monroe, R.D. Juday, "Automated self-alignment procedure for optical correlators", *Opt. Eng.* **36**, pg.1782-1791 (1997).
- [Moreno96A]** I. Moreno, V. Kober, V. Lashin, J. Campos, L.P. Yaroslavsky, M.J. Yzuel, "Color pattern recognition with circular component whitening", *Opt. Lett.* **27**, pg.498-500 (1996).
- [Moreno96B]** I. Moreno, *Diseño de sistemas multicanales en tiempo real para el reconocimiento de objetos policromáticos*, PhD. Thesis, Dept. Física, Universitat Autònoma de Barcelona (1996).
- [Moreno97]** I. Moreno, E. Ahouzi, J. Campos, M.J. Yzuel, "Real time binary amplitude phase only filters", *Appl. Opt.* **36**, pg.7428-7432 (1997).
- [Moreno98A]** I. Moreno, J.A. Davis, K.G. D'Nelly, D.B. Allison, "Transmission and phase measurement for polarization eigenvectors in twisted-nematic liquid crystal spatial light modulators", *Opt. Eng.* **37**, pg.1-5 (1998).
- [Moreno98B]** I. Moreno, J. Campos, M.J. Yzuel, V. Kober, "Implementation of bipolar real-valued input scenes in a real time optical correlator: application to color pattern recognition", *Opt. Eng.* **37**, pg.144-150 (1998).

- [Mu88]** G. Mu, X.-M. Wang, Z.-Q. Wang, "Amplitude compensated matched filter", *Appl. Opt.* **27**, pg.3461-3463 (1988).
- [Neto95]** L.G. Neto, D. Roberge, Y. Sheng, "Programmable optical phase-mostly holograms with coupled-mode modulation liquid crystal television", *Appl. Opt.* **34**, pg.1944-1950 (1995).
- [Neto96]** L.G. Neto, D. Roberge, Y. Sheng, "Full-range continuous, complex modulation by the use of two coupled-mode liquid crystal televisions", *Appl. Opt.* **35**, pg.4567-4576 (1996).
- [Nicolás2000]** J. Nicolás, M.J. Yzuel, J. Campos, "Color pattern recognition by three dimensional correlation", *Opt. Commun.* **184**, pg.335-343 (2000).
- [Nicolás2001]** J. Nicolas, C. Iemmi, J. Campos, M.J. Yzuel, "Real time correlator with liquid crystal panels: modulation optimization ", *Proc. SPIE Vol. 4419* , pg.604-607 (2001).
- [Nicolás2002A]** J. Nicolás, J. Campos, C. Iemmi, I. Moreno, M.J. Yzuel, "Convergent optical correlator alignment based on frequency filtering ", *Appl. Opt.* **41**, pg.1505-1514 (2002).
- [Nicolás2002B]** J. Nicolás, J. Campos, M.J. Yzuel, "Phase and amplitude modulation of elliptic polarization states by non-absorbing anisotropic elements: application to liquid crystal devices", *J. Opt. Soc. Am. A* **19**, pg.1013-1020 (2002).

- [Nicolás2002C]** J. Nicolás, C. Iemmi, J. Campos, M.J. Yzuel, "Optical encoding of color three-dimensional correlation", *Opt. Commun.* **209**, pg.35-43 (2002).
- [Nicolás2002D]** J. Nicolás, J. Campos, C. Iemmi, I. Moreno, M.J. Yzuel, "Filtering tests for the alignment of a convergent correlator", *Proc. SPIE Vol. 4829*, , pg.439-440 (2002).
- [Nicolás2002E]** J. Nicolás, J.Campos, M.J. Yzuel, "Elliptic light polarization modulation of liquid crystal panels", *Proc. SPIE Vol. 4829* , pg.441-442 (2002).
- [Nicolás2003]** J. Nicolás, I. Moreno, J. Campos, M.J. Yzuel, "Phase-only filtering on the three-dimensional Fourier spectrum of color images", *Appl. Opt.* **in press**, pg.- (2003).
- [O'Callaghan2001]** M.J. O'Callaghan, S.H. Perlmutter, "Serial transform optical correlator design principles", *Appl. Opt.* **40**, pg.3311-3317 (2001).
- [Partovi93]** A. Partovi, A.M. Glass, T.H. Chiu, T.H. Liu, "High-speed joint-transform optical image correlator using GaAs/AlGaAs semi-insulating multiple quantum wells and diode lasers", *Opt. Lett.* **18**, pg.906-908 (1993).
- [Pearson77]** J.J. Pearson, D.C. Hines, Jr. S. Golosman, C.D. Kuglin, "Video-rate image correlation processor", *Proc. SPIE Vol. 119* , pg.197-205 (1977).

- [Réfrégier90A]** P. Réfrégier, "Méthodes de reconnaissance des formes pour la corrélation optique", *Revue Technique THOMPSON-CSF* **22**, pg.649-734 (1990).
- [Réfrégier90B]** P. Réfrégier, "Filter design for optical pattern recognition: multicriteria optimization approach", *Opt. Lett.* **15**, pg.854-856 (1990).
- [Réfrégier91]** P. Réfrégier, "Optimal trade-off filters for noise robustness, sharpness of the correlation peak, and Horner efficiency", *Opt. Lett.* **16**, pg.829-831 (1991).
- [Réfrégier93]** P. Réfrégier, *Théorie du signal*, Masson, Paris (1993).
- [Réfrégier94]** P. Réfrégier, V. Laude, B. Javidi, "Nonlinear joint-transform correlation: an optimal solution for adaptive image discrimination and input noise robustness", *Opt. Lett.* **19**, pg.405-407 (1994).
- [Réfrégier98]** P. Réfrégier, F. Goudail, "Decision theoretical approach to nonlinear joint transform correlation", *J. Opt. Soc. Am. A* **15**, pg.61-67 (1998).
- [Reinhorn97]** S. Reinhorn, Y. Amitai, A.A. Friesem, "Compact planar optical correlator", *Opt. Lett.* **22**, pg.925-927 (1997).
- [Rollins2002]** J.M. Rollins, R.D. Juday, "Correlation techniques as applied to pose estimation in space station docking", *Proc. SPIE Vol. 4728*, O.E. Drummond, Ed., pg.218-229 (2002).

- [Romero95]** J. Romero, J.L. Nieves, A. García-Beltrán, "Human Processing of Colour Information in the Chromatic-frequency Domain", *Vision Res.* **35**, pg.867-871 (1995).
- [Rosen91]** J. Rosen, J. Shamir, "Circular harmonic phase-only filters for efficient rotation invariant pattern recognition ", *Opt. Commun.* **85**, pg.299-305 (1991).
- [Sangwine96]** S.J. Sangwine , "Fourier transforms of colour images using quaternion or hypercomplex numbers", *Electron. Lett.* **32**, pg.1979-1980 (1996).
- [Sangwine98]** S.J. Sangwine , "Colour image edge detector based on quaternion convolution ", *Electron. Lett.* **34**, pg.969-971 (1998).
- [Severcan73]** M. Severcan, *Computer Generation of Coherent Optical Filters with High Light Efficiency and Large Dynamic Range*, , PhD. Thesis, Dept. Electrical Engineering, Stanford University. (1973).
- [Soutar94]** C. Soutar, K. Lu, "Determination of the physical properties of an arbitrary twisted-nematic liquid crystal cell", *Opt. Eng.* **33**, pg.2704-2712 (1994).
- [Stolz2001]** C. Stolz, L. Bigué, P. Ambs, "Implementation of high-resolution diffractive optical elements on coupled phase and amplitude spatial light modulators", *Appl. Opt.* **40**, pg.6415-6424 (2001).

- [Thoma94]** R. Thoma, N. Hampp, "Adaptive bacteriorhodopsin-based holographic correlator for speed measurement of randomly moving three-dimensional objects", *Opt. Lett.* **19**, pg.1364-1366 (1994).
- [Thornton95]** A.L. Thornton, S.J. Sangwine, "Colour object location using complex coding in the frequency domain", *Proc. IEE 5th International conference on image processing and its applications* , pg.791-795 (1995).
- [Tudela2002]** R. Tudela, E. Martín-Badosa, I. Labastida, E. Pleguezuelos, S. Vallmitjana, I. Juvells, A. Carnicer, "Full complex Fresnel holograms displayed on liquid crystal devices", *Proc SPIE Vol. 4829* , pg.465-466 (2002).
- [Turner93]** R.M. Turner, D.A. Jared, G.D. Shark, K.M. Johnson, "Optical correlator using VLSI circuit/ferroelectric-liquid crystal electrically addressed spatial light modulators", *Appl. Opt.* **32**, pg.3094- (1993).
- [Unnikrishnan2000]** G. Unnikrishnan, M. Pohit, K. Singh, "A polarization encoded optical encryption system using ferroelectric spatial light modulator", *Opt. Commun.* **185**, pg.25-31 (2000).
- [VanderLugt64]** A. VanderLugt, "Signal detection by complex spatial filtering", *IEEE Trans. Inf. Theory* **IT-10**, pg.139-145 (1964).

- [VanderLugt67]** A. VanderLugt, "The effects of small displacements of spatial filters", *Appl. Opt.* **6**, pg.1221-1225 (1967).
- [VanderLugt92]** A. VanderLugt, *Optical Signal Processing* , Wiley, New York (1992).
- [Vijaya86]** B.V.K. Vijaya Kumar, "Minimum variance synthetic discriminant functions", *J. Opt. Soc. Am. A* **3**, pg.1579-1584 (1986).
- [Vijaya89]** B.V.K. Vijaya Kumar, Z. Bahri, "Phase-only filters with improved signal to noise ratio", *Appl. Opt.* **28**, pg.250-257 (1989).
- [Vijaya90]** B.V.K. Vijaya Kumar, L. Hasebrook, "Performance measures for correlation filters", *Appl. Opt.* **29**, pg.2997-3006 (1990).
- [Weaver66]** C.S. Weaver, J.W. Goodman, "A technique for optically convolving two functions", *Appl. Opt.* **5**, pg.1248-1249 (1966).
- [Widjaja2001]** J. Widjaja, N. Wada, Y. Ishi, W. Chujo, "Photonic packet address processor using holographic correlator", *Electron. Lett.* **37**, pg.703-704 (2001).

- [Wilkinson95]** T.D. Wilkinson, R.J. Mears, Y. Petillot, J.L. de Bougrenet de la Tocnaye, "Scale-invariant optical correlators using ferroelectric liquid-crystal spatial light modulators", *Appl. Opt.* **34**, pg.1885- (1995).
- [Yamaba93]** K. Yamaba, Y. Miyake, "Color character recognition method based on human perception", *Opt. Eng.* **32**, pg.33-40 (1993).
- [Yang82]** Y. Yang, Y.-N. Hsu, H.H. Arsenault, "Optimum circular symmetrical filters and their uses in optical pattern recognition", *Opt. Acta* **29**, pg.627-644 (1982).
- [Yaroslavsky93]** L.P. Yaroslavsky, "The theory of optimal methods for localization of objects in pictures", in *Progress in Optics Vol XXXII* , pg.145-201, E. Wolf, Ed, Elsevier, Amsterdam (1993).
- [Yaroslavsky97]** L.P. Yaroslavsky, E. Marom, "Nonlinearity optimization in nonlinear joint transform correlators", *Appl. Opt.* **36**, pg.4816- (1997).
- [Yeh99]** P. Yeh, C. Gu, *Optics of Liquid Cristal Displays* , Wiley, New York (1999).
- [Yu84]** F.T.S. Yu, "Color image recognition by spectral-spatial matched filtering", *Opt. Eng.* **23**, pg.690-695 (1984).
- [Yu86A]** F.T.S. Yu, B. Javidi, "", *Opt. Commun.* **56**, pg.384- (1986).

- [Yu86B]** F.T.S. Yu, S. Jutamulia, X.L. Huang, "Experimental application of low-cost liquid crystal TV to white-light optical signal processing", *Appl. Opt.* **25**, pg.3324-3326 (1986).
- [Yu94]** F.T.S Yu, Z. Yang, K. Pan, "Polychromatic target identification with a color liquid crystal-TV based joint transform correlator", *Appl. Opt.* **33**, pg.2170-2172 (1994).
- [Yzuel94]** M. J. Yzuel, J. Campos, "Color information in optical pattern recognition ", in *Current Trends in optics*, , pg.Academic press, London- (1994).
- [Yzuel2001]** M.J. Yzuel, J. Nicolás, I. Moreno, J. Campos, "3D filter design for color pattern recognition", *Proc. SPIE Vol. 4471*, , pg.32-42 (2001).
- [Zernike95]** F. Zernike, "How I discovered phase contrast", *Science* **121**, pg.345-349 (1995).
- [Zhou2002]** W. Zhou, M. Li, C. Wang, C. Rao, W. Jiang, "Correlation tracker for a tilt-correction adaptive optical system at the solar tower of Nanjing University", *Proc. SPIE Vol. 4926*, W. Jiang, R.K. Tyson, Eds., pg.27-39 (2002).